

4.2 that for all $p \geq 1$, the L_p -compression exponent is:

$$\alpha_p^*(\mathrm{DA}(\mathbb{T})) = \max \left\{ \frac{1}{p}, \frac{1}{2} \right\}.$$

In fact, the group $\mathrm{DA}(\mathbb{T})$ resembles very much the lamplighter *graphs* on \mathbb{T} with its affine structure introduced in Sava's thesis [Sav10a], [Sav10b]. The difference is that the moves on \mathbb{T} are somewhat twisted by the action of $\mathrm{DA}(\mathbb{T})$ on \mathbb{T} . We do not know whether the discrete affine group and Sava's lamplighter graphs are quasi-isometric, though we know the first embeds bi-Lipschitz into the second.

The main result of this paper gives an explicit description of the Poisson boundary of $\mathrm{DA}(\mathbb{T})$ with respect to any finitely supported probability measure. The Poisson boundary of a group endowed with a probability measure is a measure space giving description of the bounded harmonic functions via a Poisson formula [KV83]. An abstract description of this space is always available. However giving explicit descriptions of the Poisson boundaries for various classical groups has been an extensive subject of study over the last decades.

Kaimanovich developed two criterions called Ray and Strip criterions, which permit to describe explicitly the Poisson boundaries of, for instance, word hyperbolic groups, some Lie groups and their lattices [Kai00]. These criterions apply to many examples that had been studied before, e.g. among many others [Fur71], [Led85], [Der75] as well as [Kai91] for affine groups over the reals and [CKW94] for affine group over local fields and $\mathrm{Aff}(\mathbb{T})$. These criterions also apply for some dense subgroups in Lie groups [Bro06], [BS11].

The description of the Poisson boundary of lamplighter groups was conjectured in [KV83] to be the space of functions from the base group to the lamp group, provided that the simple random walk on the base group is transient, and shown to be trivial otherwise. It was proved for free groups of rank at least 2 by Karlsson and Woess [KW07], and for abelian groups of rank at least 5 by Erschler [Ers11], both using the Kaimanovich criterion. The Kaimanovich-Vershik conjecture was finally proved true by Lyons and Peres in [LP15]. In order to prove this conjecture, they established a new criterion which strengthened the Kaimanovich criterion. We call it the *Trap criterion*, stated below as Theorem 5.1.

In analogy with the case of lamplighter groups, we obtain the:

Theorem 1.1. *For any finitely supported non-degenerate probability measure μ , the Poisson boundary of $(\mathrm{DA}(\mathbb{T}), \mu)$ is the space of functions from \mathbb{T} to \mathfrak{S}_q , endowed with the corresponding harmonic measures.*

See Section 5 for precise statement and description of the harmonic measures. This result is perfectly analogue to lamplighter groups and Sava's lamplighter graphs. In particular as in [Sav10a], when the mean of the vertical drift is non-zero, this theorem can be proved using the Strip criterion, along the lines of [KW07]. When the mean is zero however, we use Lyons-Peres Trap criterion as in [LP15, Theorem 3.2]. An important difference here is that the well-understood affine random walk on the tree \mathbb{T} is replaced by the inverted orbit $\{o, o.w_1^{-1}, \dots, o.w_n^{-1}\}$.

This inverted orbit is a random process with memory. It is sufficiently close to a Markov process that we would be able to study it directly — see Remark 5.10. However, we rather exploit the geometric structure of the Schreier graph to get transience, and the fact that the inverted orbit respects the geometry of the tree (Fact 3.5) to build the traps — Section 5.3. An interesting feature of $\mathrm{DA}(\mathbb{T})$ is that its action on \mathbb{T} is sufficiently easy to understand the behavior of inverted orbits, though it is not the action of a group on its Cayley graph.

We also give some description of the harmonic measures in Section 5.4. In particular, we show that if two random walks have different signs of vertical drift, then their harmonic measures are mutually singular — Proposition 5.13. This result is deduced from a sublinear (logarithmic) geodesic tracking property in the tree \mathbb{T} as for instance in [Tio15].

In Section 6, we compute yet another quasi-isometric invariants related to random walks; the L_1 -isoperimetric profile and the return probability. We show that the return probability on $\text{DA}(\mathbb{T})$ has the asymptotic type

$$\mu^{(2n)}(e) \simeq e^{-n/(\log n)^2}$$

for any finitely supported symmetric measure.

The organization of this paper is the following: In Section 2, we review general properties of the full affine group $\text{Aff}(\mathbb{T})$. In Section 3, we introduce the discrete affine group $\text{DA}(\mathbb{T})$, present some basic properties of the group, and describe the metric structure of a Cayley graph. In Section 4, we give an embedding of $\text{DA}(\mathbb{T})$ into a wreath product group, and compute the L_p -compression. In Section 5, we identify the Poisson boundary of $\text{DA}(\mathbb{T})$. We establish a logarithmic geodesic tracking property in the tree \mathbb{T} , and give descriptions of the harmonic measures. In Section 6, we compute the L_1 -isoperimetric profile and the return probability of random walk with respect to any finitely supported symmetric measure.

2 The affine group of a regular tree

2.1 Generalities

We recall the background on the affine group of a regular tree. We refer to [CKW94] for a detailed exposition.

Let $\mathbb{T} = \mathbb{T}_{q+1}$ be the $q + 1$ -regular tree, equipped with the graph distance $d_{\mathbb{T}}$. An end of the tree is an equivalence class of geodesic rays (semi-infinite rays without backtracking), where two geodesic rays are equivalent when their symmetric difference is finite. We denote by $\partial\mathbb{T}$ the set of all ends. Fix, once and for all, an end ω and a distinguished vertex o in \mathbb{T} . We still denote by ω the unique geodesic ray issuing from o towards ω , and by $\omega(n)$ the vertex on ω at distance n from o . The *Busemann function* $\beta : \mathbb{T} \rightarrow \mathbb{Z}$ is defined by

$$\beta(x) = \lim_{n \rightarrow \infty} (d_{\mathbb{T}}(x, \omega(n)) - d_{\mathbb{T}}(o, \omega(n))),$$

where the limit exists for each x , since the sequence stabilizes eventually. The horocycle H_m is defined as the set of vertices whose value of β is m . Then one can regard the tree \mathbb{T} as a phylogenetic tree rooted at infinity ω , and the horocycle H_m as the m -th generation.

The group of graph automorphisms of \mathbb{T} which fix ω is called the *affine group* of \mathbb{T} and denoted by $\text{Aff}(\mathbb{T})$. (Changing the end ω yields an isomorphic group as the tree is regular.) Observe that each affine transformation sends one horocycle to the other and this fact motivates to define a map from the group $\text{Aff}(\mathbb{T})$ to the integers as follows: For each $\gamma \in \text{Aff}(\mathbb{T})$ and $x \in \mathbb{T}$, we define $\Phi(\gamma) := \beta(x \cdot \gamma) - \beta(x)$. Note that $\Phi(\gamma)$ does not depend on the choice of x , and thus the map $\Phi : \text{Aff}(\mathbb{T}) \rightarrow \mathbb{Z}$ is, in fact, a homomorphism. Define the subgroup $\text{Hor}(\mathbb{T})$ of $\text{Aff}(\mathbb{T})$ as the kernel of Φ . The group $\text{Hor}(\mathbb{T})$ preserves each horocycle.

One can verify that for the topology induced by the pointwise convergence on \mathbb{T} , the group $\text{Hor}(\mathbb{T})$ is an inductive limit of compact groups (the subgroups fixing a vertex on the geodesic ray ω). Moreover, the exact sequence

$$1 \rightarrow \text{Hor}(\mathbb{T}) \rightarrow \text{Aff}(\mathbb{T}) \rightarrow \mathbb{Z} \rightarrow 1$$

splits. Indeed, choosing some α such that $\Phi(\alpha) = 1$ determines the semidirect product decomposition of $\text{Aff}(\mathbb{T})$ by $\text{Hor}(\mathbb{T}) \rtimes \mathbb{Z}$. The group $\text{Aff}(\mathbb{T})$ is a locally compact topological group which is non-unimodular by [SW90] and amenable as a topological group by [Neb88].




Figure 1: The affine 3-regular tree with its labelling.

2.2 Labelling of the tree

Let us introduce a labelling of the tree \mathbb{T} . We identify the horocycle $H_m = \beta^{-1}(m)$ with the set of all sequences indexed by $] -\infty, m] \cap \mathbb{Z} = \llbracket -\infty, m \rrbracket$, taking values in the alphabet $\mathcal{A}_q = \{0, \dots, q-1\}$, and constant equal to 0 on a subinterval $\llbracket -\infty, n \rrbracket$. In the phylogenic description, the k -ancestor of a sequence is obtained by deleting the last (rightmost) k entries. The edges of the tree link each vertex to its first ancestor.

Up to changing the labelling, we can choose $\omega(n)$ to be the constant sequence equal to 0 on $\llbracket -\infty, -n \rrbracket$, and α to be the shift $(u_k)_{k \leq m} \mapsto (u_{k-1})_{k \leq m+1}$ from H_m to H_{m+1} .

The labelling extends naturally to the boundary, parametrized by the set of sequences indexed by \mathbb{Z} taking values in \mathcal{A}_q and constant equal to 0 on a neighborhood of $-\infty$, plus the empty sequence corresponding to our chosen end ω — see Figure 1. We warn that the sign of index n for $\omega(n)$ and H_n is opposite.

2.3 Notion of portrait

If an element γ of $\text{Hor}(\mathbb{T})$ fixes a vertex v , then it induces an automorphism of the rooted subtree \mathbb{T}_v obtained by restriction to the root vertex v and all its descendants. Automorphism groups of rooted trees have been largely studied — see for instance [Gri00], [Nek05], [Zuk08] and references therein. The notion of portrait is a useful tool in their study, and can be generalized here as follows.

Consider the symmetric group \mathfrak{S}_q acting by permutations on \mathcal{A}_q . Let $\gamma : \mathbb{T} \rightarrow \mathfrak{S}_q$ be a function denoted $v \mapsto \gamma[v]$. Assume that $\gamma[\omega(n)] = e$ for all but finitely many n . Such a function defines an element of $\text{Hor}(\mathbb{T})$, still denoted γ , as follows.

For $n \geq N = \max\{n : \gamma[\omega(n)] \neq e\}$, the action of γ on the vertex $\omega(n)$ is trivial. Now assume that we know the action of γ on a vertex v identified with its labelling sequence. The action of γ on the descendant vt , where $t \in \mathcal{A}_q$, is given recursively by the “automata rule”:

$$vt.\gamma = (v.\gamma)(t.\gamma[v]), \quad (1)$$

where $t.\gamma[v]$ is the action of $\gamma[v] \in \mathfrak{S}_q$ on $t \in \mathcal{A}_q$. This permits to define by induction the action of γ on all \mathbb{T} since any vertex v has the form $\omega(n)t_1 \dots t_k$ for some $n \geq N$ and t_i in \mathcal{A}_q .

By construction, any element $\gamma \in \text{Hor}(\mathbb{T})$ can be obtained bijectively from such a function $\mathbb{T} \rightarrow \mathfrak{S}_q$ still denoted by γ and called its *portrait*. The composition rule for the portrait of a product is:

$$\gamma\gamma'[v] = \gamma[v]\gamma'[v.\gamma]. \quad (2)$$

We use right action, meaning γ is applied first and then γ' . We refer to [Gri00] for details in the case of a rooted tree.

By the semidirect product decomposition $\text{Aff}(\mathbb{T}) = \text{Hor}(\mathbb{T}) \rtimes \mathbb{Z}$, an element of the affine group is identified with a pair $g = (\gamma, m) = \gamma\alpha^m$. The product rule is:

$$gh = (\gamma, m)(\gamma', m') = (\gamma(\gamma')^{\alpha^m}, m + m'),$$

where the conjugate $\gamma^\alpha = \alpha\gamma\alpha^{-1}$ has the portrait $\gamma^\alpha[v] = \gamma[v.\alpha]$.

By abuse of notation, we denote by $g[v]$ instead of $\gamma[v]$ when $g = \gamma\alpha^m$, so for instance $\alpha[v] = e$ for all v . The combination of the previous line with (2) gives: for $g, h \in \text{Aff}(\mathbb{T})$, $v \in \mathbb{T}$,

$$gh[v] = g[v]h[v.g]. \quad (3)$$

It follows that the portrait of a product gh is obtained from that of g after multiplying by $h[u]$ at position $u.g^{-1}$, for every u in \mathbb{T} .

3 The discrete affine group

3.1 Generalities

Since the subgroup of $\text{Hor}(\mathbb{T})$ consisting of elements with finitely supported portrait is invariant under conjugation by the shift, the following is a subgroup of $\text{Aff}(\mathbb{T})$, which we call the *discrete affine group*.

$$\text{DA}(\mathbb{T}) := \{g \in \text{Aff}(\mathbb{T}) : g[v] = e \text{ for all but finitely many } v\}.$$

This is neither a lattice of the affine group $\text{Aff}(\mathbb{T})$ (which cannot exist by the non-unimodularity), nor a discrete subgroup of the topological affine group, but a finitely generated dense subgroup of $\text{Aff}(\mathbb{T})$ — see Proposition 3.1.

For each element $\sigma \in \mathfrak{S}_q$ and each vertex $v \in \mathbb{T}$, denote δ_v^σ the horocyclic element with portrait given by

$$\delta_v^\sigma[w] = \begin{cases} e & \text{if } w \neq v \\ \sigma & \text{if } w = v. \end{cases}$$

For each $v \in \mathbb{T}$, the morphism $\sigma \mapsto \delta_v^\sigma$ is injective of finite image $\delta_v^{\mathfrak{S}_q} \subset \text{Hor}(\mathbb{T})$ isomorphic to \mathfrak{S}_q .

Proposition 3.1. *The group $\text{DA}(\mathbb{T})$*

1. *is generated by the finite set $S_0 = \{\alpha, \alpha^{-1}\} \cup \delta_o^{\mathfrak{S}_q}$,*
2. *is dense in $\text{Aff}(\mathbb{T})$ in the pointwise convergence topology,*
3. *is elementary amenable,*
4. *does not satisfy any group law.*

Recall that a group G satisfies a *group law* if there exists a non-trivial irreducible word w in an alphabet of size k such that $w(g_1, \dots, g_k) = \text{id}$ for all $g_i \in G$, $1 \leq i \leq k$.

Proof. (1) As for all $m \in \mathbb{Z}$, we have $(\delta_o^\sigma)^{\alpha^m} = \delta_{\omega(m)}^\sigma$, it is sufficient to check that $\delta_{\omega(m)}^{\mathfrak{S}_q}$ for m in \mathbb{Z} generates $\text{Hor}(\mathbb{T}) \cap \text{DA}(\mathbb{T})$. It follows from the following folklore lemma applied to show that $\delta_{\omega(m)}^{\mathfrak{S}_q}$ for $m = -M$ to M generate the subgroup of $\text{Hor}(\mathbb{T})$ consisting of elements with portrait trivial outside the q -ary subtree rooted at $\omega(M)$ of depth $2M + 1$ (which is naturally isomorphic to the group of automorphism of this finite rooted tree).

Lemma 3.2. *Let $T(k)$ be a q -regular rooted tree of finite depth $k + 1$, with a distinguished ray from the root to a leaf. For $0 \leq i \leq k$, denote $\Delta_i \simeq S_q$ the subgroup of automorphisms of $T(k)$ with portrait supported on the i -th vertex of the ray. Then the union $\cup_{i=0}^k \Delta_i$ is a generating set of $\text{Aut}(T(k))$.*

Proof. We use induction on k and the permutational wreath product isomorphism $\text{Aut}(T(k)) \simeq \text{Aut}(T(k-1)) \wr_{\{0, \dots, q-1\}} \Delta_0$. By induction, $\Delta_1, \dots, \Delta_k$ generate the first copy of $\text{Aut}(T(k-1))$. The other copies are obtained by conjugation by Δ_0 . \square

(2) We have to prove that given an element $g = (\gamma, m)$ in $\text{Aff}(\mathbb{T})$ and a finite collection of points in the tree, we can find an element in $\text{DA}(\mathbb{T})$ with the same action as γ on these points. For this, choose $\omega(n)$ a common ancestor of all the given points. Any element γ' in $\text{Hor}(\mathbb{T})$ fixing $\omega(n)$ and with the same portrait as γ on a finite subtree containing $\omega(n)$ and the given points will have the same action as the horocyclic part of g . Thus $\gamma' \alpha^m$ is the required element.

(3) This group is a cyclic extension of the subgroup of $\text{Hor}(\mathbb{T})$ the elements of which have finite support. The latter is locally finite, whence the group $\text{DA}(\mathbb{T})$ is elementary amenable.

(4) is obtained from the following theorem by Abért.

Theorem 3.3 (Theorem 1 in [Abé05]). *Let G be a group acting on a set X , satisfying the following separability condition: for every finite subset Y of X , the pointwise stabilizer*

$$G_Y := \bigcap_{v \in Y} \{g \in G : gv = v\}$$

does not stabilize any point outside Y . Then G does not satisfy any group law.

We observe that the action of $\text{DA}(\mathbb{T})$ on $\partial\mathbb{T} \setminus \{\omega\}$ satisfies the separability condition in Theorem 3.3. Indeed, it follows from the fact that for every $v \in \mathbb{T}$, we have $\delta_v^{\mathfrak{S}_q} \subset \text{DA}(\mathbb{T})$. \square

Remark 3.4. We can replace \mathfrak{S}_q by any of its subgroup S with transitive action on $\mathcal{A}_q = \{0, \dots, q-1\}$, and obtain a group $\text{DA}_S(\mathbb{T})$ finitely generated, elementary amenable and dense in $\text{Aff}_S(\mathbb{T})$ in which all permutations of the portrait are in S rather than arbitrary.

3.2 Description of the word metric

Our first aim here is to understand how to compute the portrait of a given word in the generators $S_0 = \{\alpha, \alpha^{-1}\} \cup \delta_o^{\mathfrak{S}_q}$ of the discrete affine group $\text{DA}(\mathbb{T})$.

By induction, let g be such a word and h be a generator. By (3), the portrait of gh satisfies that for $v \in \mathbb{T}$,

$$gh[v] = g[v]h[v.g] = \begin{cases} g[v] & \text{if } h \in \{\alpha, \alpha^{-1}\} \\ g[v]\delta_o^\sigma[v.g] = g[v]\delta_{o.g^{-1}}^\sigma[v] & \text{if } h = \delta_o^\sigma. \end{cases} \quad (4)$$

In particular, the portrait is the same as that of g except possibly at the point $o.g^{-1}$. This makes it important to understand what happens to this point when we multiply by a generator in S_0 . So we want to express $o.(gh)^{-1} = o.h^{-1}g^{-1}$ in terms of $o.g^{-1}$ and the generator h . We record the

Fact 3.5. *We have the following multiplication rules:*

1. If $h = \delta_o^\sigma$, then $o.(\delta_o^\sigma)^{-1} = o$, so $o.(g\delta_o^\sigma)^{-1} = o.g^{-1}$.
2. If $h = \alpha$, then $o.\alpha^{-1} = \omega(1)$ is the first ancestor of o , so $o.(g\alpha)^{-1}$ is the ancestor of $o.g^{-1}$.
3. If $h = \alpha^{-1}$, then $o.\alpha = o0$, so by the automata rule (1)

$$o.(g\alpha^{-1})^{-1} = o0.g^{-1} = (o.g^{-1})0.g^{-1}[o],$$

which is a first descendant of $o.g^{-1}$ determined by $g^{-1}[o] = (g[o.g^{-1}])^{-1}$.

Consider a word $w_n = x_1 \dots x_n$ (which may not be reduced) in the generators S_0 and denote by $w_i = x_1 \dots x_i$ its prefix of length i . It follows from (4) that the support of the portrait of w_n is included in the inverted orbit $O_n = \{o, o.w_1^{-1}, o.w_2^{-1}, \dots, o.w_n^{-1}\}$.

The inverted orbit O_n is a sequence of points in the Schreier graph $\mathcal{G}(\text{DA}(\mathbb{T}), S_0, o)$ — see Figure 2. Usually, it is extremely difficult to understand the inverted orbit of a word in a given action. In the present particular case, Fact 3.5 permits to describe clearly O_n from the word w_n . We make use of it to estimate the word metric in $\text{DA}(\mathbb{T})$.

The set $S_1 = \delta_o^{\mathfrak{S}_q} \{\alpha^{\pm 1}\} \delta_o^{\mathfrak{S}_q}$ is also a generating set of $\text{DA}(\mathbb{T})$, as follows easily from Proposition 3.1(1). It can be compared to the “switch-walk-switch” generating sets in lamplighter groups, for which the word metric is given by travelling salesman paths on the base graph. This is also the case here.

In a graph G , given a starting vertex x , an end vertex y and a collection A of vertices, we denote by $\text{TSP}_G(A; x, y)$ the length of the shortest path starting at x , visiting all vertices in A and ending in y , called a travelling salesman path.

Let us denote by $\text{supp}(g) = \{v \in \mathbb{T} : g[v] \neq e\}$ the support of the portrait of an element g of $\text{DA}(\mathbb{T})$. We set $\ell(g)$ to be the number of edges of the smallest subtree $T(g)$ containing $\text{supp}(g) \cup \{o, o.g^{-1}\}$.

Proposition 3.6. *The word metric in $\text{DA}(\mathbb{T})$ with respect to the generating set $S_1 = \delta_o^{\mathfrak{S}_q} \{\alpha^{\pm 1}\} \delta_o^{\mathfrak{S}_q}$ is given by*

$$|g|_{S_1} = \text{TSP}_{\mathbb{T}}(\text{supp}(g); o, o.g^{-1}) = 2\ell(g) - d_{\mathbb{T}}(o, o.g^{-1}).$$

for all $g \in \text{DA}(\mathbb{T}) \setminus \delta_o^{\mathfrak{S}_q}$. Moreover $|\delta_o^\sigma|_{S_1} = 2$ for σ non-trivial in \mathfrak{S}_q .

The second equality is obtained by the well-known description of travelling salesman paths in trees. We could also express the word metric with respect to S_0 but it would be slightly less elegant.

Proof. By Fact 3.5, the inverted orbit of a word in the generators S_1 is a path in the tree \mathbb{T} . By (4) when we multiply by a generator in S_1 , we modify the portrait at the two endpoints of the corresponding edge in the path. So the length of a representative word of g is at least the length of a path solution to a travelling salesman problem in \mathbb{T} starting in o ending in $o.g^{-1}$ and visiting all vertices in $\text{supp}(g)$. This gives the lower bound.

There remains to give a representative word of g of right length. For this, let us first describe a path solution to the travelling salesman problem. The edges of this path are precisely those of the tree $T(g)$. The edges located on the geodesic from o to $o.g^{-1}$ are covered exactly once, and those not on this geodesic are covered exactly twice. We can define a word w_n in S_1 with inverted orbit precisely this path, and such that the portrait of the prefix w_i coincides with the portrait of g restricted to the vertices that have been visited for the last time before time i .

Indeed, assume w_i is given, we can obtain w_{i+1} as follows.

If time i is the last visit of the path to vertex $o.w_i^{-1}$, multiply by an element in $\delta_o^{\mathfrak{S}_q}$ that ensures $w_{i+1}[o.w_i^{-1}] = g[o.w_i^{-1}]$. Then multiply by α or α^{-1} to reach the next point of the path.




Figure 2: Schreier graph $\mathcal{G}(\text{DA}(\mathbb{T}), S_0, o)$ for $q = 3$.

If time i is not the last visit of the path to vertex $o.w_i^{-1}$, multiply by an element in $\delta_o^{\mathfrak{S}_q}$ that ensures $o.w_{i+1}^{-1}$ is the next vertex of the path (this is possible by Fact 3.5). Then multiply by α or α^{-1} to reach the next point of the path.

At final time n , multiply by an element in $\delta_o^{\mathfrak{S}_q}$ that ensures $w_n[o.w_n^{-1}] = g[o.w_n^{-1}]$. \square

3.3 The Schreier graph of the action on \mathbb{T}

For future use, we describe the *Schreier graph* associated with the action of $\text{DA}(\mathbb{T})$ on \mathbb{T} with respect to the generating set S_0 of Proposition 3.1 and the orbit of o . We denote it by $\mathcal{G} = \mathcal{G}(\text{DA}(\mathbb{T}), S_0, o)$. The set of vertices is $\mathbb{T} = \{o.x : x \in \text{DA}(\mathbb{T})\}$, and the set of directed (labelled) edges is defined by $\{o.x, o.xs\}$ for some $s \in S_0$. Both $\{o.x, o.xs\}$ and $\{o.xs, o.x\}$ are edges by the symmetry of S_0 , and there are also loops. We draw a picture of \mathcal{G} without loops and multiple edges in Figure 2. We denote the graph distance by $d_{\mathcal{G}}$.

Lemma 3.7. *For every $x \in \text{DA}(\mathbb{T})$, we have*

$$d_{\mathbb{T}}(o, o.x) \leq d_{\mathcal{G}}(o, o.x) \leq 2d_{\mathbb{T}}(o, o.x).$$

Proof. First note that in \mathbb{T} we have $d_{\mathbb{T}}(o, o.x) = d_{\mathbb{T}}(o, o.x\sigma)$ for all $\sigma \in \delta_o^{\mathfrak{S}_q}$. For each $o.x \in \mathcal{G}$, let w be a reduced word which realizes a directed path from o to $o.x$ with $o.x = o.w$. The number of $\{\alpha, \alpha^{-1}\}$ in w bounds the number of steps from o to $o.x$ in \mathbb{T} ; this gives $d_{\mathbb{T}}(o, o.x) \leq d_{\mathcal{G}}(o, o.x)$.

On the other hand, first let us consider those vertex $o.x$ in the subtree \mathbb{T}_o . For a (unique) geodesic path from o to $o.x$, each step in order to obtain $o.x$ we may apply a $\sigma \in \delta_o^{\mathfrak{S}_q}$; hence 2 times the number of steps gives the bound of $d_{\mathcal{G}}(o, o.x)$. In general, we note that a geodesic path from o and $o.x$ in \mathbb{T} is realized in such a way that there exists a subtree $\mathbb{T}_{o.\alpha^{-l}}$ for some $l \geq 0$,

the path is a concatenation of the segment between o and $o.\alpha^{-l}$, and the geodesic path between $o.\alpha^{-l}$ and $o.x$. The geodesic between $o.\alpha^{-l}$ and $o.x$ is the isomorphic image of the geodesic between o and $o.x\alpha^l$ via α^{-l} in \mathbb{T} ; applying the argument used in above special case, we have $d_G(o, o.x) \leq 2d_{\mathbb{T}}(o, o.x)$. \square

4 Embeddings into wreath products and L_p -compression

4.1 Embeddings into wreath products

We consider the group $\text{DA}(\mathbb{T})$ and fix the set of generators S_1 as in Proposition 3.6, and write the corresponding word metric by $d_{\text{DA}}(x, y) := |x^{-1}y|_{\text{DA}}$, where we do not write it such as $|x^{-1}y|_{S_1}$ for clarity of notation. We compare this metric in $\text{DA}(\mathbb{T})$ with a group of wreath product.

Given two groups H and F , recall that the wreath product $H \wr F$ is defined as $\oplus_F H \rtimes F$, where the semidirect product is given by the left action: $\varphi \mapsto \varphi(x^{-1}\cdot)$ for $x \in F$ and $\varphi \in \oplus_F H$.

The set $\{(\delta_e^h, e)\}_{h \in \mathfrak{S}_q} \cup \{(\mathbf{1}, s)\}_{s \in S_F}$, where e is the identity element of the base group F , δ_e^h is the element in $\oplus_F H$ given by h at e and the identity otherwise, and $\mathbf{1}$ is the identity everywhere on F , generates $H \wr F$ and is often called the “switch-or-walk” generating set. The set

$$\{(\delta_e^{h_1}, e)(\mathbf{1}, s)(\delta_e^{h_2}, e) : h_1, h_2 \in H, s \in S_F\}, \quad (5)$$

is another generating set of $H \wr F$ called “switch-walk-switch” generating set, more convenient for our purpose.

Let us consider the wreath product where $H = \mathfrak{S}_q$ and $F = F_{q+1}$ is the free product of $q+1$ copies of \mathbb{Z}_2 . The Cayley graph of F_{q+1} with respect to the standard symmetric set of generators S_F is a $q+1$ -regular tree. We fix an identification of this tree with \mathbb{T} . We consider the Cayley graph of $\Gamma := \mathfrak{S}_q \wr F_{q+1}$ associated with the set of generators in (5), and denote the corresponding word metric by $d_\Gamma(x, y) := |x^{-1}y|_\Gamma$.

Lemma 4.1. *There exists an isometric embedding $f : \text{DA}(\mathbb{T}) \rightarrow \Gamma = \mathfrak{S}_q \wr F_{q+1}$, i.e. for all x, y in $\text{DA}(\mathbb{T})$, one has $d_\Gamma(f(x), f(y)) = d_{\text{DA}}(x, y)$.*

Proof. We define the map $f : \text{DA}(\mathbb{T}) \rightarrow \Gamma = \mathfrak{S}_q \wr F_{q+1}$ by

$$g \mapsto (\{g[v]\}_{v \in \mathbb{T}}, o.g^{-1})$$

under the identification F_{q+1} with \mathbb{T} . It is well-known that the word metric d_Γ of wreath product with switch-walk-switch generating set is given by

$$|(\varphi, x)|_\Gamma = \begin{cases} \text{TSP}(\text{supp}\varphi; e, x), & \text{if } (\varphi, x) \neq (\delta_e^h, e) \\ 2 & \text{if } (\varphi, x) = (\delta_e^h, e) \text{ with } h \neq e \end{cases}$$

where $\text{TSP}(\text{supp}\varphi; e, x)$ denotes the minimal length of path in \mathbb{T} from e to x visiting all the points in $\text{supp}\varphi$. This is an isometry by Proposition 3.6. \square

Remark 4.2. The argument shows that there are bi-Lipschitz embeddings of $\text{DA}(\mathbb{T})$ into Sava’s lamplighter graphs $\mathfrak{S}_q \wr \mathbb{T}$ and into the wreath product group $\mathfrak{S}_q \wr G$ as soon as the group G contains an infinite bi-Lipschitz embedded binary tree.

Of course, the group $\text{DA}(\mathbb{T})$ is not quasi-isometric to Γ because the latter is non-amenable. It would be interesting to show that it is not quasi-isometric to Sava’s lamplighter graphs, nor to $\mathfrak{S}_q \wr (\mathbb{Z}_2 \wr \mathbb{Z})$. It might be subtle as their quasi-isometry classes cannot be distinguished by isoperimetric profile, nor by L_p -compression.

4.2 L_p -compression of the discrete affine group

Recall that for a finitely generated group Γ , and for $1 \leq p < \infty$, the L_p -compression exponent $\alpha_p^*(\Gamma)$ is the supremum of all those $\alpha \geq 0$ for which there exists a Lipschitz map $\Gamma \rightarrow L_p$ such that for some constant $c > 0$ and for all x, y in Γ ,

$$\|f(x) - f(y)\|_p \geq cd_\Gamma(x, y)^\alpha,$$

where d_Γ denotes a word metric in Γ . If $p = 2$, then $\alpha_2^*(\Gamma)$ is called the *Hilbert compression exponent*. These are quasi-isometric group invariants introduced in [GK04]; for backgrounds and related results, see e.g. [NP08], [NP11] and references therein. For all $1 \leq p < \infty$, we determine the exact values of L_p -compression exponents for $\text{DA}(\mathbb{T})$.

Proposition 4.3. *The group $\text{DA}(\mathbb{T})$ admits a bi-Lipschitz embedding into L_1 . Moreover, for $1 \leq p < \infty$,*

$$\alpha_p^*(\text{DA}(\mathbb{T})) = \max\left\{\frac{1}{p}, \frac{1}{2}\right\}.$$

The proposition follows from several known results: the upper bound is obtained by the method using escape rate of random walks due to Naor and Peres [NP11], and the lower bound is derived from the corresponding result on wreath products over free groups by Cornulier, Stalder and Valette [CSV12].

Proof of Proposition 4.3. First we show the upper bound for $\alpha_p^*(\text{DA}(\mathbb{T}))$. For a finitely generated group G , let $\beta^*(G)$ be the supremum of all those $\beta \geq 0$ for which there exists a finite symmetric set of generators S , and a constant $c > 0$ such that for every $n \geq 1$,

$$\mathbb{E}[d_G(e, w_n)] \geq cn^\beta,$$

where $\{w_n\}_{n=0}^\infty$ is the simple random walk on the Cayley graph of G associated with S , starting at the identity e . Naor and Peres show that if G is amenable, then

$$\alpha_p^*(G) \leq \max\left\{\frac{1}{p}, \frac{1}{2}\right\} \frac{1}{\beta^*(G)}, \quad (6)$$

see [NP11, (5) in Section 1 and Theorem 1.3]. In $\text{DA}(\mathbb{T})$, the simple random walk on any Cayley graph has a linear rate of escape since the Poisson boundary is non-trivial (Theorem 5.3), and thus we have $\beta^*(\text{DA}(\mathbb{T})) = 1$; then (6) gives the desired upper bound for $\alpha_p^*(\text{DA}(\mathbb{T}))$.

Next we show the lower bound for $\alpha_p^*(\text{DA}(\mathbb{T}))$. A result of Cornulier, Stalder and Valette implies that $\alpha_1^*(\mathfrak{S}_q \wr F_{q+1}) = 1$; in fact, the group admits a bi-Lipschitz embedding into L_1 [CSV12, Proof of Proposition 7.2]. Combining with Lemma 4.1, we deduce that $\text{DA}(\mathbb{T})$ admits a bi-Lipschitz embedding into L_1 , in particular, $\alpha_1^*(\text{DA}(\mathbb{T})) = 1$. In general, we have

$$\alpha_p^*(\text{DA}(\mathbb{T})) \geq \max\left\{\frac{1}{p}, \frac{1}{2}\right\} \alpha_1^*(\text{DA}(\mathbb{T}))$$

by [NP11, p.103]. This yields the required lower bound for $\alpha_p^*(\text{DA}(\mathbb{T}))$. \square

5 The Poisson boundary of the discrete affine group

Let μ be a probability measure on $\text{DA}(\mathbb{T})$. We consider the associated right random walk $w_n = x_1 \dots x_n$, where the factors $x_i = \gamma_i \alpha^{m_i}$ form a sequence of independent random variables of law μ . We will always assume that μ is *non-degenerate*, i.e. its support generates $\text{DA}(\mathbb{T})$ as a semigroup.

By Fact 3.5, understanding the random inverted orbit $O_n = \{o, o.w_1^{-1}, \dots, o.w_n^{-1}\}$ is important to describe the behavior of the random walk. For a fixed n , the random word $w_n^{-1} = x_n^{-1} \dots x_1^{-1} = \tilde{w}_n$ coincides with the *left* random walk of law $\tilde{\mu}$ given by $\tilde{\mu}(g) = \mu(g^{-1})$.

Both right and left random walks on the topological affine group of \mathbb{T} have been studied by Cartwright, Kaimanovich and Woess [CKW94]. Note that as we use right actions on the tree, the roles of left and right random walks here are unfortunately exchanged with their paper.

As already observed in [CKW94], the inverted orbit $\{o.\tilde{w}_n\}$ is not a Markov process on \mathbb{T} , except in extremely particular case — see Remark 5.4. On the contrary, the direct orbit $\{o.w_n\}$ is always a Markov chain, because the transitions depend only on space and not on time.

5.1 Poisson boundary

We recall here basic facts about Poisson boundaries. We refer to [KV83], [Kai00], [Fur02] or [LP16, Chapter 14] for more informations. The space of trajectories $\Omega := \Gamma^{\mathbb{N}}$ of a random walk of law μ on a group Γ is endowed with the (left) diagonal action of the group Γ via each coordinate and the probability measure $\mathbb{P}_x = \pi_* \mu^{\mathbb{N}}$ obtained by pushing forward the Bernoulli measure $\mu^{\mathbb{N}}$ on the space of increments by $\pi : (x_1, x_2, \dots) \mapsto (w_0, w_1, w_2, \dots)$ where $w_0 = x$ corresponds to the starting point. Moreover the shift naturally acts on the space of trajectories by $\tau : (w_1, w_2, \dots) \mapsto (w_2, w_3, \dots)$. Note that the shift τ commutes with the group action. Let \mathcal{I} be the σ -field of shift-invariant events. The *Poisson boundary* of (Γ, μ) is the measurable space (Ω, \mathcal{I}) endowed with the family of probability measures $\{\mathbb{P}_x\}_{x \in \Gamma}$. Notice that arbitrary two measures \mathbb{P}_x and \mathbb{P}_y are mutually absolutely continuous since μ is non-degenerate; it follows that $L^\infty(\Omega, \mathcal{I}, \mathbb{P}_x)$ and $L^\infty(\Omega, \mathcal{I}, \mathbb{P}_y)$ are naturally isomorphic. Let us denote by $\mathbb{P} := \mathbb{P}_e$.

We call a function $f : \Gamma \rightarrow \mathbb{R}$ is μ -harmonic if $\sum_{h \in \Gamma} f(gh)\mu(h) = f(g)$ for all g in Γ . Let $H^\infty(\Gamma, \mu)$ be the space of all bounded μ -harmonic functions on Γ endowed with ℓ^∞ -norm. One can check that the linear map from $L^\infty(\Omega, \mathcal{I}, \mathbb{P})$ to $H^\infty(\Gamma, \mu)$ defined by

$$\varphi \mapsto \int_{\Omega} \varphi(\omega) d\mathbb{P}_x(\omega),$$

(where the right hand side is a function of x in Γ), gives an isometric isomorphism as Banach spaces by the bounded martingale convergence theorem.

In practice, many explicit description of Poisson boundaries, e.g. [KW07], [Bro06], [BS11], [Sav10b], have been done using the “Strip criterion” due to Kaimanovich [Kai00]. Indeed, the Strip criterion also works for a special random walk on $\text{DA}(\mathbb{T})$ (where one can adapt the methods in [KW07]). In order to deal with a general random walk, we use the following enhanced version of this criterion.

Theorem 5.1 (Lyons-Peres Trap criterion [LP15] Corollary 2.3, and [LP16] Proposition 14.42). *Let \mathcal{A} be a Γ -invariant sub- σ -field of \mathcal{I} . Assume that for every $\varepsilon > 0$, there exists a random sequence of finite subsets $\{D_\ell\}_{\ell=1}^\infty$ of Γ and a constant $c > 0$ such that*

- (i) *each D_ℓ is a (set-valued) measurable function with respect to \mathcal{A} ,*
- (ii) *almost surely there exists N such that $|D_\ell| \leq e^{\varepsilon \ell}$ for all $\ell \geq N$,*
- (iii) *$\limsup_{\ell \rightarrow \infty} \mathbb{P}(\exists n \geq \ell, w_n \in D_\ell) \geq c$.*

then $\mathcal{A} = \mathcal{I}$ modulo \mathbb{P} -null sets. In particular, (Ω, \mathcal{A}) gives a realization of the Poisson boundary.

The sets D_ℓ can be thought of as traps which can be defined in terms of \mathcal{A} by (i), are small by (ii) and catch the random walk with positive probability by (iii).

5.2 Description of the Poisson boundary: Main result

Our aim here is to describe the Poisson boundary of $\text{DA}(\mathbb{T})$. Let us first assume that the step distribution μ is finitely supported. Denote by R the minimal integer such that for all $g \in \text{supp}(\mu)$,

$$|\Phi(g)| \leq R, \quad \text{and} \quad g[v] = e \quad \text{when} \quad d_{\mathbb{T}}(v, o) > R. \quad (7)$$

We assume that μ is non-degenerate. In many cases we also assume that μ is *aperiodic*, i.e. for any given group elements g and g' there exists a number $l \geq 1$ such that the support of the l -times convolution power of μ contains both g and g' . If μ is not aperiodic, then one may replace it by $\tilde{\mu} := (1/2)\mu + (1/2)\delta_e$, which is aperiodic and non-degenerate; moreover μ -harmonic functions are always $\tilde{\mu}$ -harmonic functions and the converse is also true.

Cartwright, Kaimanovich and Woess observed that the sequence $\{o.\tilde{w}_n\}$ converges to an end $\partial\mathbb{T}$ almost surely. More precisely:

Proposition 5.2 (A special case of Theorem 2 in [CKW94]). *Assume that μ is non-degenerate on $\text{DA}(\mathbb{T})$, and consider the standard compactified topology in $\mathbb{T} \cup \partial\mathbb{T}$.*

- (i) *If $\mathbb{E}_{\tilde{\mu}}\Phi > 0$, then $\{o.\tilde{w}_n\}$ converges to a random element in $\partial\mathbb{T} \setminus \{\omega\}$ almost surely.*
- (ii) *If $\mathbb{E}_{\tilde{\mu}}\Phi \leq 0$, then $o.\tilde{w}_n \rightarrow \omega$ almost surely.*

As the group $\text{DA}(\mathbb{T})$ acts by automorphisms of the tree \mathbb{T} , we always have $d_{\mathbb{T}}(o, v) = d_{\mathbb{T}}(o.\tilde{w}_n, v.\tilde{w}_n)$. Combined with (3) and (7), It follows by (3) and (7) that for a fixed v in \mathbb{T} the sequence $w_n[v]$ changes values only when $d_{\mathbb{T}}(o.\tilde{w}_n, v) \leq R$, so is eventually constant by Proposition 5.2. Denote $\gamma_{\infty}[v] \in \mathfrak{S}_q$ its limit. This means that the sequence of portraits converges almost surely pointwise to a final configuration given by a Borel measurable map

$$\gamma_{\infty} : \Omega \rightarrow \prod_{\mathbb{T}} \mathfrak{S}_q.$$

Let $\mathcal{A} = \sigma(\gamma_{\infty})$ be the sub- σ -field of \mathcal{I} generated by final configurations. The map γ_{∞} is Γ -equivariant with respect to a natural Γ -action on $\prod_{\mathbb{T}} \mathfrak{S}_q$. We will show that the hypothesis of Lyons-Peres Trap criterion stated in Theorem 5.1 apply to \mathcal{A} and deduce the following more precise version of Theorem 1.1.

Theorem 5.3. *Assume that μ is a finitely supported, non-degenerate probability measure on $\text{DA}(\mathbb{T})$. Then the space $\prod_{\mathbb{T}} \mathfrak{S}_q$ of final configurations with the distribution $\gamma_{\infty*}\mathbb{P}$ is a realization of the Poisson boundary of $(\text{DA}(\mathbb{T}), \mu)$.*

This statement is completely analogous of what happens for lamplighter groups. The Trap criterion was designed in [LP15] to show that final configurations realize the Poisson boundary for random walks in lamplighter groups.

Remark 5.4. In her thesis [Sav10a, Section 5.3], Sava introduced some lamplighter random walks on the graph $\mathbb{Z}_2 \wr \mathbb{T}$ and identified their Poisson boundaries as spaces of lamp configurations. Random walks in the group $\text{DA}(\mathbb{T})$ resemble very much random walks in the graph $\mathbb{Z}_2 \wr \mathbb{T}$, as the portrait corresponds to the lamp configuration.

Sava's results were improved by Lyons and Peres [LP15, Theorem 3.2]. Our proof will follow their ideas. A major difference in our case is that the random trajectory $\{o.\tilde{w}\}_n$ induced on \mathbb{T} is usually *not* a Markov process. It has some memory encoded in the portrait.

A particular case where this process is actually Markov is obtained when μ is the equidistribution measure on the set $S_1 = \delta_o^{\mathfrak{S}_q} \{\alpha^{\pm 1}\} \delta_o^{\mathfrak{S}_q}$. Indeed we see by Fact 3.5 that the entry $w_n[o.\tilde{w}_n]$ is completely randomized before each move by α^{-1} . This is no longer true for general step distribution, even when μ is symmetric, i.e., $\mu = \tilde{\mu}$.

5.3 Proof of Theorems 1.1 and 5.3

The measure μ on $\text{DA}(\mathbb{T})$ induces a random walk on \mathcal{G} given by the forward orbit $\{o.w_n\}$. This is a Markov chain on \mathcal{G} with transition probability $p(x, y) = \mu(\{s : x.s = y\})$ for $x, y \in \mathcal{G}$.

Lemma 5.5. *For μ on $\text{DA}(\mathbb{T})$ as in Theorem 5.3, the random walk induced by μ on the Schreier graph $\mathcal{G} = \mathcal{G}(\text{DA}(\mathbb{T}), S_0, o)$ is transient.*

In particular, if $B_{\mathcal{G}}(o, R)$ is the ball of radius R centered at o in \mathcal{G} , there exist n_0 and $c_R > 0$ such that

$$\mathbb{P}(\forall n \geq n_0, o.w_n \notin B_{\mathcal{G}}(o, R)) \geq c_R > 0.$$

Proof. We regard the Schreier graph \mathcal{G} as a network endowed with conductance 1 on each edge. We use reduction of the network by contracting edges corresponding to $\sigma \in \delta_o^{\oplus q}$, and observe that the reduced network contains a copy of \mathbb{T}_o , which is transient. The Rayleigh monotonicity implies that the simple random walk on \mathcal{G} is transient as well (e.g. [LP16, Chapter 2]). For the network endowed with transition probability $p(x, y)$, which is uniformly bounded away from 0, but need not be reversible, the random walk $\{o.w_n\}$ is still transient by [Woe00, Theorem 2.25 and Theorem 3.1]. Here n_0 is the smallest number of steps to get out from the ball $B_{\mathcal{G}}(o, R)$. \square

Assume $\mathbb{E}_{\tilde{\mu}}\Phi > 0$ (resp. $\mathbb{E}_{\tilde{\mu}}\Phi \leq 0$), let us define τ_{ℓ} for $\ell > 0$ as the first time when the random trajectory $o.\tilde{w}_n$ hits or crosses the horocycle H_{ℓ} (resp. $H_{-\ell}$). This coincides with the first time $\Phi(w_n)$ hits or crosses ℓ (resp. $-\ell$), thus is almost surely well defined.

Removing the point $o.\tilde{w}_n$ splits \mathbb{T} into $q + 1$ subtrees. We denote T_{ℓ} the subtree containing o and all other previous points of the inverted orbit. The following lemma asserts that the random trajectory $\{o.\tilde{w}_n\}$ does not come back to T_{ℓ} with positive probability. This is a quantitative form of transience.

Lemma 5.6. *There exists a constant c depending only on μ such that for all $\ell > 0$,*

$$\mathbb{P}(\forall n \geq \tau_{\ell}, o.\tilde{w}_n \notin T_{\ell} \mid \mathcal{F}_{\tau_{\ell}}) \geq c > 0,$$

where $\mathcal{F}_{\tau_{\ell}}$ is the σ -algebra generated by sequence $\{\tilde{w}_i\}_{i \leq \tau_{\ell}}$ up to τ_{ℓ} . In particular, for all $\ell > 0$,

$$\mathbb{P}(\forall n \geq \tau_{\ell}, o.\tilde{w}_n \notin T_{\ell}) \geq c > 0.$$

This lemma is obvious when $\mathbb{E}_{\tilde{\mu}}\Phi \neq 0$ for in this case we can use transience of the projected random walk $\beta(o.\tilde{w}_n) = -\Phi(w_n)$ on the integers, which is finitely supported and drifted.

Proof of Lemma 5.6. Recall that τ_{ℓ} is the first time when the random trajectory $o.\tilde{w}_n$ hits or crosses the horocycle $H_{-\ell}$. Then $\{\tilde{w}_{\tau_{\ell} + n}\}_{n \geq 0}$ has the same distribution as $\{w_n\}_{n \geq 0}$ by the strong Markov property. Let R be as in (7). If $d_{\mathcal{G}}(o, o.\tilde{w}_{\tau_{\ell} + n}) \geq 2R$, then $d_{\mathbb{T}}(o, o.\tilde{w}_{\tau_{\ell} + n}) \geq R$ by Lemma 3.7, and so $d_{\mathbb{T}}(o.\tilde{w}_{\tau_{\ell}}, o.\tilde{w}_{\tau_{\ell} + n}) \geq R$. Therefore if $o.w_n \notin B_{\mathcal{G}}(o, 2R)$ for all $n \geq \tau_{\ell}$, then $o.\tilde{w}_n \notin B_{\mathbb{T}}(o, R)$ for all $n \geq \tau_{\ell}$.

In fact, the last condition implies that $o.\tilde{w}_n \notin T_{\ell}$ for all $n \geq \tau_{\ell}$. Indeed, by definition of R , the condition implies that either all $o.\tilde{w}_n$ for $n \geq \tau_{\ell}$ are in T_{ℓ} , which cannot happen because the projection $\{\Phi(w_n)\}$ is recurrent on integers, or none.

By Lemma 5.5, we obtain

$$\mathbb{P}(\forall n \geq \tau_{\ell} + n_0, o.\tilde{w}_n \notin T_{\ell} \mid \mathcal{F}_{\tau_{\ell}}) \geq \mathbb{P}(\forall n \geq n_0, o.w_n \notin B_{\mathcal{G}}(o, 2R)) \geq c_{2R} > 0. \quad (8)$$

To get Lemma 5.6, observe that for any $\ell' \geq \ell$, the probability that $o.\tilde{w}_n$ does not belong to T_{ℓ} for $\tau_{\ell} \leq n \leq \tau_{\ell'}$ is bounded below by a constant $c_1 > 0$. Choosing $\ell' = n_0(R + 1) + \ell$ guarantees that $o.\tilde{w}_n$ does not belong to T_{ℓ} for $\tau_{\ell'} \leq n \leq \tau_{\ell'} + n_0$. Lemma 5.6 holds with $c = c_1 c_{2R} > 0$. \square

Lemma 5.7. *Assume that μ is aperiodic. The limit $\lim o.\tilde{w}_n \in \partial\mathbb{T}$ is almost surely the unique point of accumulation of $\text{supp}(\gamma_\infty)$.*

Proof. We give the proof in the case $\mathbb{E}_\mu\Phi \leq 0$. Since μ is non-degenerate and aperiodic, one can find an integer j such that $\mu^{\otimes j}(v)$ and $\mu^{\otimes j}(v')$ are both positive, where $v = \alpha^{R+1}$ and $v' = \sigma\alpha^{R+1}$ with $\sigma \in \delta_o^{\mathbb{S}^q}$ non trivial (in the case $\mathbb{E}_\mu\Phi > 0$, take negative powers of α instead) and R defined in (7).

There almost surely exist infinitely many integers n such that the random word w_n ends with v or v' at $n = \tau_\ell$, where ℓ is a positive multiple of $R + 1$. Moreover, by Lemma 5.6, almost surely there is an infinite subsequence $(\ell_k)_{k \geq 0}$ of this set of integers such that

$$o.\tilde{w}_n \notin T_{\ell_k} \text{ for all } n \geq \tau_{\ell_k}. \quad (9)$$

Indeed, for each ℓ , the probability that (9) does not hold is at most $1 - c$, so the probability that this does not happen for n successive cases is at most $(1 - c)^n$. This shows that $\text{supp}(\gamma_\infty)$ has almost surely at most one accumulation point. There remains to show that $\text{supp}(\gamma_\infty)$ is almost surely infinitely supported.

Now for each k , at least one of the two possibilities v or v' guarantees to have a non-trivial portrait at $o.\tilde{w}_{\tau_{\ell_k}}$. This also holds for γ_∞ by (9) and this can be done independently for different k as the ℓ_k differ by at least $R + 1$. Note that v' appears infinitely many times, because the probabilities of occurrence of v or v' are given by independent Bernoulli laws of the same non-trivial parameter. So the size of $\text{supp}(\gamma_\infty)$ is almost surely infinite. \square

Proof of Theorems 1.1 and 5.3. In order to apply Theorem 5.1, define the random sequence of subsets $\{D_\ell\}_{\ell=1}^\infty$ of $\text{DA}(\mathbb{T})$ as follows. Let $c_1 > 0$ be a parameter to be fixed below. We denote $o_{c_1\ell}$ the point at distance $c_1\ell$ from o on the unique geodesic ray from o to $\lim o.\tilde{w}_n \in \partial\mathbb{T}$. Let $T_{c_1\ell}$ be the subtree of $\mathbb{T} \setminus \{o_{c_1\ell}\}$ containing o .

The subset D_ℓ consists of all elements $g = (\gamma, m)$ such that $c_1\ell \leq m \leq c_1\ell + R$ and

$$\gamma[v] = \begin{cases} \gamma_\infty[v] & \text{if } d_{\mathbb{T}}(v, o_{c_1\ell}) > R \text{ and } v \in T_{c_1\ell}, \\ e & \text{if } d_{\mathbb{T}}(v, o_{c_1\ell}) > R \text{ and } v \notin T_{c_1\ell}. \end{cases}$$

Clearly the size of D_ℓ is bounded by a constant depending only on q and R , so point (ii) is satisfied. By Lemma 5.7, the limit $\lim o.\tilde{w}_n$ is almost surely the unique point of accumulation of $\text{supp}(\gamma_\infty)$, so the set D_ℓ is measurable with respect to $\mathcal{A} = \sigma(\gamma_\infty)$ and point (i) is satisfied.

Now let $\tau_{c_1\ell}$ as above be the first hitting or crossing time of $H_{c_1\ell}$ when $\mathbb{E}_\mu\Phi > 0$ (resp. $H_{-c_1\ell}$ when $\mathbb{E}_\mu\Phi \leq 0$). Also denote $o'_{c_1\ell}$ the point at the intersection of the horocycle $H_{c_1\ell}$ (or $H_{-c_1\ell}$) and the geodesic between o and $o.\tilde{w}_{\tau_{c_1\ell}}$. Lemma 5.6 asserts that with probability greater than $c > 0$, we have $o'_{c_1\ell} = o_{c_1\ell}$ and $w_{\tau_{c_1\ell}}$ belongs to D_ℓ . So point (iii) is satisfied when $c_1 > R$ because $\tau_{c_1\ell} \geq c_1\ell/R$ by (7).

Note that when $\mathbb{E}_\mu\Phi \leq 0$, we always have $o'_{c_1\ell} = o_{c_1\ell} = \omega(c_1\ell)$. \square

Remark 5.8. Note that the existence of the subsequence (ℓ_k) in the proof of Lemma 5.7 provides another proof of Proposition 5.2, where the end is given by the intersection of all $\mathbb{T} \setminus T_{\ell_k}$. The original proof used a general result on transience in non-unimodular groups by Guivarc'h, Keane and Roynette: as μ is non-degenerate and $\text{DA}(\mathbb{T})$ is dense in $\text{Aff}(\mathbb{T})$ by Proposition 3.1, the closed subgroup generated by the support of μ is $\text{Aff}(\mathbb{T})$ which is non-unimodular. It follows by [GKR77, Theorem 51] that the random walks $\{w_n\}$ and $\{\tilde{w}_n\}$ are transient in $\text{Aff}(\mathbb{T})$, i.e. they leave every compact set almost surely; for details, see [CKW94, Theorem 2].

Remark 5.9. In the case $\mathbb{E}_\mu\Phi \neq 0$, Theorem 5.3 still holds when μ has a finite first moment. The proof above can be adapted as follows. Given $\varepsilon > 0$, the sets D_ℓ now consist of all elements such

that $c_1\ell \leq m \leq c_1\ell + \varepsilon\ell$ and

$$\gamma[v] = \begin{cases} \gamma_\infty[v] & \text{if } d_{\mathbb{T}}(v, o_{c_1\ell}) > \varepsilon\ell \text{ and } v \in T_{c_1\ell}, \\ e & \text{if } d_{\mathbb{T}}(v, o_{c_1\ell}) > \varepsilon\ell \text{ and } v \notin T_{c_1\ell}. \end{cases}$$

The drift of the random walk induced on the integers still gives Lemma 5.6 and point (i). The first moment condition ensures that the sequence of increments almost surely has a sublinear word norm. Combined with Proposition 3.6, this shows that (7) can be replaced by the almost sure existence of N such that for all $n \geq N$,

$$|\Phi(x_n)| \leq \varepsilon n \quad \text{and} \quad x_n[v] = e \text{ when } d_{\mathbb{T}}(v, o) > \varepsilon n.$$

The drift of the random walk induced on the integers still gives Lemma 5.6 and so $w_{\tau_{c_1\ell}}$ still belongs to D_ℓ with positive probability. Point (i) also still holds.

Finally, the law of large numbers gives a constant $c_2 > 0$ depending only on μ such that $\tau_\ell/\ell \rightarrow c_2$ almost surely. So point (iii) is satisfied when $c_1 > 1/c_2$.

Remark 5.10. We sketch here another proof of Lemma 5.6 when $\mathbb{E}_\mu \Phi = 0$, considering only the random trajectory $o.\tilde{w}_n$ in the tree \mathbb{T} . Looking at times when $\Phi(w_n)$ first visits negative integers, we see that the line $\omega = o.\alpha^{\mathbb{Z}}$ is recurrent. The metric projection $\text{proj}(o.\tilde{w}_n)$ onto this line is a random process on \mathbb{Z} with memory encoded in the portrait of w_n . However it is stochastically dominated by a random walk on \mathbb{Z} drifted towards $-\infty$. The reason is that at times when $o.\tilde{w}_n$ is not on the line, the projection may only decrease, and at times when it is on the line, the projection may decrease by $x < 0$ with probability $p(x) = \Phi_*\mu(x)$ but it may increase by $x > 0$ with probability $p(x)$ satisfying only $\sum_{x>0} xp(x) \leq \sum_{x>0} x\Phi_*\mu(x) - c_1$ for some $c_1 > 0$ independent of the past. A precise coupling argument can be developed, but we prefer to omit the tedious details.

Remark 5.11. The Schreier graph \mathcal{G} shares some visual similarities with the Schreier graph of Thompson's group F acting on dyadic rationals pictured in [Sav15]. Both are essentially infinite binary rooted trees with infinitely many rays attached to it. Transience of this Schreier graph was used by Kaimanovich to show that the Poisson boundary of F with respect to finitely supported measures is non-trivial [Kai16]. In the discrete affine group, we can actually identify the Poisson boundary because the inverted orbit respects the geometric structure of \mathbb{T} , and this permits to define the traps. It is not known whether the μ -boundary of F described in [Kai16] is the whole Poisson boundary. Note that this boundary is actually trivial for some infinitely supported measures [JZ16].

5.4 Description of the harmonic measure

The aim here is to give a description of the distribution (the harmonic measure) $\gamma_{\infty*}\mathbb{P}$ on the space $\Pi_{\mathbb{T}}\mathfrak{S}_q$ of final configurations, which is the Poisson boundary by Theorem 5.3. We first describe a sublinear geodesic tracking property in the tree \mathbb{T} when $\mathbb{E}_{\tilde{\mu}}\Phi \neq 0$, in the spirit of Tiozzo [Tio15].

Proposition 5.12. *Let μ be as in Section 5.2. If $\mathbb{E}_{\tilde{\mu}}\Phi \neq 0$, then there exists a constant $C > 0$ such that for \mathbb{P} -almost every sample, there is a (random) geodesic ray ξ from o converging to a point $\partial\mathbb{T}$ satisfying that*

$$\limsup_{n \rightarrow \infty} \frac{d_{\mathbb{T}}(o.\tilde{w}_n, \xi)}{\log n} \leq C.$$

In the case when $\mathbb{E}_{\tilde{\mu}}\Phi < 0$, the geodesic ray ξ is a deterministic one converging to ω , while in the case when $\mathbb{E}_{\tilde{\mu}}\Phi > 0$, the geodesic ray ξ is a random one converging to some point in $\partial\mathbb{T} \setminus \{\omega\}$.

Applying a general theorem by Tiozzo [Tio15, Theorem 6] immediately gives deviation of order $o(n)$ instead of $\log n$ for $\mathbb{E}_{\tilde{\mu}}\Phi \neq 0$.

Proof. First we prove the claim when $\mathbb{E}_{\tilde{\mu}}\Phi > 0$. Let $F := \max_{g \in \text{supp}\mu} d_{\mathbb{T}}(o, o.g)$. We define a sequence of regeneration time; namely, let

$$\bar{\tau}(n) := \inf\{k > 0 : \forall i \geq k, \Phi(\check{w}_i) > Fn\}$$

for $n \geq 0$. We have $\bar{\tau}(n) < \bar{\tau}(n+1)$ for all n . Note that $\bar{\tau}(n)$ are not stopping times. Given the sequence $\{\bar{\tau}(n) : n \geq 0\}$, a family of excursions $\{X_i^{(n)} : i \in I_n\}$ for $I_n := [0, \bar{\tau}(n+1) - \bar{\tau}(n))$ defined by

$$X_i^{(n)} := \Phi(\check{w}_{i+\bar{\tau}(n)}) - \Phi(\check{w}_{\bar{\tau}(n)})$$

are independent, and each excursion is an asymmetric random walk on \mathbb{Z} drifted toward $+\infty$ conditioned on the event $X_{\bar{\tau}(n+1)-\bar{\tau}(n)-1} \in (-\infty, F]$. By comparison with the asymmetric random walk on \mathbb{Z} , we have that there exist constants $c_1, c_2 > 0$ such that for every $n \geq 0$ and for every $M \geq 0$,

$$\mathbb{P}\left(\max_{i \in I_n} X_i^{(n)} \geq M \mid \{\bar{\tau}(m) : m \geq 0\}\right) \leq c_1 e^{-c_2 M}.$$

Therefore given the sequence $\{\bar{\tau}(n) : n \geq 0\}$, there exists a constant $C > 0$ such that

$$\max_{k \in [\bar{\tau}(n), \bar{\tau}(n+1))} d_{\mathbb{T}}(o, \check{w}_k) \leq C \log n,$$

for all large enough n almost surely. Notice that the F -neighbourhood $\mathbb{T}(n, F)$ of the subtree \mathbb{T}_{v_n} where $v_n := o.\check{w}_{\bar{\tau}(n)}$ are nested: $\mathbb{T}(n, F) \supset \mathbb{T}(n+1, F) \supset \dots$, and thus there exists a geodesic ray ξ from o converging to some ξ in $\partial\mathbb{T} \setminus \{\omega\}$, where we use the same symbol ξ , such that $d_{\mathbb{T}}(o, \check{w}_{\bar{\tau}(n)}, \xi) \leq F$. Clearly, we have $n \leq \bar{\tau}(n)$. We have that \mathbb{P} -almost surely

$$d_{\mathbb{T}}(o, \check{w}_n, \gamma) \leq C \log n + F$$

for all large enough n , and obtain the claim.

Next we show the claim when $\mathbb{E}_{\tilde{\mu}}\Phi < 0$. We define the sequence of stopping times: for $n \geq 0$,

$$\tau(n) := \min\{k \geq 0 : \Phi(\check{w}_k) \leq -Fn\}.$$

The family of excursions $\{X_i^{(n)} : i \in I_n\}$ for $n = 0, 1, 2, \dots$ defined in a similar way to above are independent, and each excursion is an asymmetric random walk on \mathbb{Z} drifted toward $-\infty$ up to a random stopping time. Again comparison with the walk on \mathbb{Z} yields the proof. \square

Let us denote the harmonic measure by ν_{μ} which is $\gamma_{\infty*}\mathbb{P}$ on the space $\Pi_{\mathbb{T}}\mathfrak{S}_q$. We show that ν_{μ} have distinct supports depending on the signs of $\mathbb{E}_{\tilde{\mu}}\Phi$.

Proposition 5.13. *Let ν_- , ν_0 and ν_+ be the harmonic measures for aperiodic step distributions μ_- , μ_0 , and μ_+ respectively satisfying $\mathbb{E}_{\tilde{\mu}_-}\Phi < 0$, $\mathbb{E}_{\tilde{\mu}_0}\Phi = 0$ and $\mathbb{E}_{\tilde{\mu}_+}\Phi > 0$. Then ν_- , ν_0 and ν_+ have distinct supports in $\Pi_{\mathbb{T}}\mathfrak{S}_q$. In particular, they are mutually singular.*

Proof. First we observe that, by Lemma 5.7, the support of final configurations $\text{supp}\gamma_{\infty}$ has the accumulation point ω in the case when $\mathbb{E}_{\tilde{\mu}}\Phi \leq 0$, and a (random) accumulation point in $\partial\mathbb{T} \setminus \{\omega\}$ in the case when $\mathbb{E}_{\tilde{\mu}}\Phi > 0$. This shows that the support of ν_+ is distinct from that of ν_- and ν_0 . There remains to show that ν_- and ν_0 have distinct supports.

Let us consider the infinite subtree $\mathbb{T}_{\ell} = \mathbb{T}_{\omega(\ell)}$ rooted at $\omega(\ell)$ and its finite subtree $\mathbb{T}_{\ell, f(\ell)}$ consisting of all descendants up to distance $f(\ell)$ from the root $\omega(\ell)$. We define a subset in \mathbb{T} by

$$D(\ell, f) := \mathbb{T}_{\ell} \cup \bigcup_{m \geq \ell} \mathbb{T}_{m, f(m)},$$

Let us define a set of configuration $\hat{D}(f) := \{\varphi \in \Pi_{\mathbb{T}}\mathfrak{S}_q : \exists \ell, \text{supp}\varphi \subset D(\ell, f)\}$. Proposition 5.12 gives a constant $C > 0$ such that $\nu_-(\hat{D}(C \log)) = 1$.

There remains to show that $\nu_0(\hat{D}(C \log)) = 0$. In fact, for $f(\ell) = \ell + g(\ell)$ with $g(\ell) \leq \ell$, we claim that $\nu_0(\hat{D}(f)) = 0$.

We argue as in the proof of Lemma 5.7, where v and v' were defined. By properties of zero mean random walks on \mathbb{Z} , there are almost surely infinitely many ℓ_k 's such that the trajectory $o.\tilde{w}_n$ reaches level $g(\ell_k) + 1$ in the tree \mathbb{T}_{ℓ_k} . Let τ_k be the stopping time when $o.\tilde{w}_n$ first comes back to a level $\leq g(\ell_k)$. We consider only the almost surely infinitely many k 's where w_{τ_k} ends with v or v' . By Lemma 5.6, there is a fixed positive probability that the trajectory never visits again the subtree above vertex $o.\tilde{w}_{\tau_k}$. At least one of the two possible endpoints v or v' guarantees that the final configuration is non trivial in $\mathbb{T}_{\ell_k} \setminus \mathbb{T}_{\ell_k, f(\ell_k)}$, proving the claim. \square

Remark 5.14. Considering only the accumulation point of Lemma 5.7 gives a μ -boundary of the random walk (see [Kai00],[Fur02] for a definition) which is a quotient of the Poisson boundary. In topological (spread-out measures) random walks on $\text{Aff}(\mathbb{T})$, this accumulation point is actually the Poisson boundary by [CKW94].

6 Return probability and isoperimetric profile

For a finitely generated group Γ and for a symmetric probability measure μ on the group Γ , the L_1 -isoperimetric profile $\Lambda_{1,\Gamma,\mu} : [1, \infty) \rightarrow \mathbb{R}$ is defined by

$$\Lambda_{1,\Gamma,\mu}(v) := \inf \left\{ \frac{1}{2} \sum_{x,y \in \Gamma} |f(x) - f(y)| \mu(y) : |\text{supp } f| \leq v, \|f\|_1 = 1 \right\}.$$

By using the co-area formula, the L_1 -isoperimetric profile $\Lambda_{1,\Gamma,\mu}$ is equivalent to

$$\Lambda_{1,\Gamma,\mu}(v) \simeq \inf \left\{ \frac{1}{|U|} \sum_{x,y \in \Gamma} 1_U(x) 1_{\Gamma \setminus U}(xy) \mu(y) : |U| \leq v \right\}, \quad (10)$$

where given functions $f, g : (0, \infty) \rightarrow (0, \infty)$ (where the domain might be restricted on integers), we write $f \lesssim g$ if there exist constants $C_1, C_2 > 0$ such that $f(t) \leq C_1 g(C_2 t)$ for all large enough t , and write $f \simeq g$ if we have both $f \lesssim g$ and $f \gtrsim g$.

For a finite symmetric set of generators S of a group Γ , let μ_S be the uniform measure on S . For another finite symmetric set of generators H of Γ , we have $\Lambda_{1,\Gamma,\mu_S} \simeq \Lambda_{1,\Gamma,\mu_H}$. In fact, one can check that the equivalence class (the *asymptotic type*) of Λ_{1,Γ,μ_S} is a quasi-isometric invariant of the group Γ .

The L_1 -isoperimetric profile gives an estimate for the return probability $\mu^{(2n)}(e)$, where $\mu^{(2n)}$ denotes the $2n$ -times convolution power of μ . Again, for any two finite symmetric set of generators S, H of a group Γ , we have $\mu_S^{(2n)}(e) \simeq \mu_H^{(2n)}(e)$, and in fact, the asymptotic type of $\mu_S^{(2n)}(e)$ is a quasi-isometric invariant of the group Γ as shown by Pittet and Saloff-Coste [PSC00, Theorem 1.2].

Proposition 6.1. *Let μ be a finitely supported symmetric probability measure on $\text{DA}(\mathbb{T})$. Assume that the support of μ generates the whole group as a semigroup. Then*

$$\Lambda_{1,\text{DA}(\mathbb{T}),\mu}(v) \simeq \frac{1}{\log \log v}, \quad (11)$$

and

$$\mu^{(2n)}(e) \simeq e^{-n/(\log n)^2}. \quad (12)$$

Proof. Given $r \in \mathbb{N}$, consider the set L_r which consists of vertices in the horocycle H_r that are descendants of o , that is those vertices with labeling $\dots 000.w$, $w \in \{0, \dots, q-1\}^r$. Let C_r be the

subgroup of elements of $\text{Hor}(\mathbb{T})$ such that the support of the portrait is contained in L_r ,

$$C_r = \{\gamma \in \text{Hor}(\mathbb{T}) : \text{supp}\gamma \subseteq L_r\}.$$

Then C_r is isomorphic to the product $\mathfrak{S}_q^{|L_r|}$. Consider the transition kernel that chooses one coordinate uniformly from L_r and update the permutation to be uniform in \mathfrak{S}_q , that is

$$\eta_r = \frac{1}{|L_r|} \sum_{v \in L_r} \frac{1}{|\mathfrak{S}_q|} \sum_{\sigma \in \mathfrak{S}_q} \delta_v^\sigma,$$

where δ_v^σ is the indicator function on the element γ with portrait σ at v and trivial everywhere else. We apply Erschler's isoperimetric inequality as in [SCZ15a, Proposition 3.1], then we have

$$\Lambda_{1,C_r,\eta_r}(v) \geq \frac{1}{2K} \text{ for all } v \leq \frac{1}{K} |\mathfrak{S}_q|^{|L_r|/K}$$

for some absolute constant $K > 0$. Here for elements in the support of η_r , the word distance relative to S_0 in $\text{DA}(\mathbb{T})$ is bounded by $2r + 1$. Therefore by comparison, for the uniform measure μ_{S_0} on the set of generators S_0 of $\text{DA}(\mathbb{T})$, we have

$$\Lambda_{1,\text{DA}(\mathbb{T}),\mu_{S_0}}(v) \geq \frac{1}{2Cr} \text{ for all } v \leq \frac{1}{K} |\mathfrak{S}_q|^{|L_r|/K} = \frac{1}{K} (q!)^{q^r/K}.$$

This gives

$$\Lambda_{1,\text{DA}(\mathbb{T}),\mu_{S_0}}(v) \gtrsim 1/\log \log v. \quad (13)$$

On the other hand, for $r \in \mathbb{N}$, consider the finite subset of $\text{DA}(\mathbb{T})$,

$$U := \{\gamma \alpha^i : 0 \leq i \leq r, \text{supp}\gamma \subseteq L'_r\},$$

where L'_r is the set of vertices descendant of o belonging to $\cup_{0 \leq s \leq r} H_s$. Then for μ_{S_0} , we have

$$\Lambda_{1,\text{DA}(\mathbb{T}),\mu_{S_0}}(v) \leq \frac{2}{|S_0|r} \text{ for all } v \geq |U| = r(q!)^{q^r},$$

and this yields $\Lambda_{1,\text{DA}(\mathbb{T}),\mu_{S_0}}(v) \lesssim 1/\log \log v$. As we noted, $\Lambda_{1,\text{DA}(\mathbb{T}),\mu_{S_0}} \simeq \Lambda_{1,\text{DA}(\mathbb{T}),\mu}$ for a general finitely supported probability measure μ that generates the whole group as a semigroup.

The lower bound (13) of $\Lambda_{1,\text{DA}(\mathbb{T}),\mu}$ gives an upper bound of the return probability:

$$\mu^{(2n)}(e) \lesssim e^{-n/(\log n)^2},$$

by using the Nash inequality [Cou96] (see also [SCZ15b]); we omit the detail. For the return probability lower bound, we can use a simple argument that on the event that the projection to \mathbb{Z} is confined in $[-r, r]$, then the portrait is confined to $B_{\mathbb{T}}(o, r)$. There exist constants $c_1, c_2 > 0$ such that for all large enough r and n , one has

$$\mathbb{P} \left(\max_{0 \leq k \leq n} |\Phi(\check{w}_k)| \leq r \right) \geq c_1 e^{-c_2 n/r^2}$$

[Ale92, Lemma 1.2]. Then

$$\mu^{(2n)}(e) \geq \frac{1}{2r(q!)^{|B_{\mathbb{T}}(o,r)|}} e^{-Cn/r^2}.$$

Optimise choice of r we obtain a matching lower bound. \square

Acknowledgements. J.B. is partially supported by project AGIRA ANR-16-CE40-0022. R.T. is supported by JSPS Grant-in-Aid for Young Scientists (B) (No. 26800029).

References

- [Abé05] M. Abért. Group laws and free subgroups in topological groups. *Bulletin of the London Mathematical Society*, 37(04):525–534, 2005.
- [Ale92] G. Alexopoulos. A lower estimate for central probabilities on polycyclic groups. *Can. J. Math.*, 44(5):897–910, 1992.
- [Bro04] S. Brofferio. Renewal theory on the affine group of an oriented tree. *Journal of Theoretical Probability*, 17(4):819–859, 2004.
- [Bro06] S. Brofferio. The Poisson boundary of random rational affinities. *Annales de l’institut Fourier*, 56(2):499–515, 2006.
- [BS11] S. Brofferio and B. Schapira. Poisson boundary of $GL_d(\mathbb{Q})$. *Israel Journal of Mathematics*, 185(1):125–140, 2011.
- [CKW94] D. I. Cartwright, V. A. Kaimanovich, and W. Woess. Random walks on the affine group of local fields and of homogeneous trees. *Ann. Inst. Fourier, Grenoble*, 44(4):1243–1288, 1994.
- [Cou96] T. Coulhon. Ultracontractivity and Nash type inequality. *Journal of Functional Analysis*, 141:510–539, 1996.
- [CSV12] Y. Cornuier, Y. Stalder, and A. Valette. Proper actions of wreath products and generalizations. *Transactions of the American Mathematical Society*, 364(6):3159–3184, 2012.
- [Der75] Y. Derriennic. Marche aléatoire sur le groupe libre et frontière de Martin. *Z. Wahrscheinlichkeitstheor. Verw. Geb.*, 32:261–276, 1975.
- [Ers11] A. Erschler. Poisson-Furstenberg boundary of random walks on wreath products and free metabelian groups. *Comment. Math. Helv.*, 86:113–143, 2011.
- [Fur71] H. Furstenberg. Random walks and discrete subgroups of Lie groups. *Advances in probability and related topics*, 1:1–63, 1971.
- [Fur02] A. Furman. Random walks on groups and random transformations. In North-Holland, editor, *Handbook of dynamical systems, Vol. 1a chap.12 Hasselblatt and Katok*, pages 931–1014, 2002.
- [GK04] E. Guentner and J. Kaminker. Exactness and uniform embeddability of discrete groups. *J. London Math. Soc.*, 70(3):703–718, 2004.
- [GKR77] Y. Guivarc’h, M. Keane, and B. Roynette. *Marches Aléatoires sur les Groupes de Lie, Lecture Notes in Math.*, 624. Springer Berlin-Heidelberg-New York, 1977.
- [Gri00] R. Grigorchuk. Just infinite branch groups. In *New Horizons in pro-p-Groups*, volume 184, pages 121–179. Progress in Mathematics, 2000.
- [JZ16] K. Juschenko and T. Zheng. Infinitely supported Liouville measures of Schreier graphs. arXiv:1608.03554, 2016.
- [Kai91] V. A. Kaimanovich. Poisson boundaries of random walks on discrete solvable groups. In *Probability measures on groups X (Oberwolfach 1990)*, pages 205–238, New York, 1991. Penum.
- [Kai00] V. A. Kaimanovich. The Poisson formula for groups with hyperbolic properties. *Ann. of Math. (2)*, 152(3):659–692, 2000.
- [Kai16] V. A. Kaimanovich. Thompson’s group F is not Liouville. arXiv:1602.02971, 2016.

- [KV83] V. A. Kaimanovich and A. Vershik. Random walks on discrete groups : boundary and entropy. *Annals of Probability*, 11(3):457–490, 1983.
- [KW07] A. Karlsson and W. Woess. The Poisson boundary of lamplighter random walks on trees. *Geom. Dedicata*, 124:95–107, 2007.
- [Led85] F. Ledrappier. Poisson boundaries of discrete groups of matrices. *Israel Journal of Mathematics*, 50:319–336, 1985.
- [LP15] R. Lyons and Y. Peres. Poisson boundaries of lamplighter groups: proof of the Kaimanovich-Vershik conjecture. *arXiv:1508.01845*, 2015.
- [LP16] R. Lyons and Y. Peres. *Probability on Trees and Networks*. Cambridge University Press, 2016.
- [Neb88] C. Nebbia. Amenability and Kunze-Stein property for groups acting on a tree. *Pacific Journal of Mathematics*, 135(2):371–380, 1988.
- [Nek05] V. Nekrashevych. *Self-similar groups*. Amer. Math. Soc., 2005.
- [NP08] A. Naor and Y. Peres. Embeddings of discrete groups and the speed of random walks. *International Mathematics Research Notices*, 2008:rnn076, 2008.
- [NP11] A. Naor and Y. Peres. L_p compression, traveling salesmen, and stable walks. *Duke Math. J.*, 157(1):53–108, 2011.
- [PSC00] C. Pittet and L. Saloff-Coste. On the stability of the behavior of random walks on groups. *The Journal of Geometric Analysis*, 10(4):713–737, 2000.
- [Sav10a] E. Sava. *Lamplighter Random Walks and Entropy-Sensitivity of Languages*. Ph.D. thesis. Technische Universität Graz, 2010.
- [Sav10b] E. Sava. A note on the Poisson boundary of lamplighter random walks. *Monatshefte für Mathematik*, 159(4):379–396, 2010.
- [Sav15] D. Savchuk. Schreier graphs of actions of Thompson’s group F on the unit interval and on the cantor set. *Geom. Dedicata*, 175:355–372, 2015.
- [SCZ15a] L. Saloff-Coste and T. Zheng. Isoperimetric profiles and random walks on some permutation wreath products. *arXiv:1510.08830*, 2015.
- [SCZ15b] L. Saloff-Coste and T. Zheng. Random walks and isoperimetric profiles under moment conditions. *arXiv:1501.05929*, 2015.
- [SW90] P. Soardi and W. Woess. Amenability, unimodularity, and the spectral radius of random walks on infinite graphs. *Mathematische Zeitschrift*, 205:471–486, 1990.
- [Tio15] G. Tiozzo. Sublinear deviation between geodesics and sample paths. *Duke Math. J.*, 164(3):511–539, 2015.
- [Woe00] W. Woess. *Random walks on infinite graphs and groups*, volume 138 of *Cambridge Tracts in Mathematics*. Cambridge University Press, 2000.
- [Zuk08] A. Zuk. Groupes engendrés par des automates. *Séminaire Bourbaki, Astérisque*, 311:141–174, 2008.

JÉRÉMIE BRIEUSSEL — UNIVERSITÉ DE MONTPELLIER — jeremie.brieussel@umontpellier.fr

RYOKICHI TANAKA — TOHOKU UNIVERSITY — rtanaka@m.tohoku.ac.jp

TIANYI ZHENG — UC SAN DIEGO — tzheng2@math.ucsd.edu